

# Physician's Rx



## Compression Levels:

Medical:  20-30 mmHg     30-40 mmHg     40-50 mmHg

## Style:

calf <input type="checkbox"/> L <input type="checkbox"/> R	thigh <input type="checkbox"/> L <input type="checkbox"/> R	thigh w/ waist att. <input type="checkbox"/> L <input type="checkbox"/> R	panty <input type="checkbox"/>	maternity panty <input type="checkbox"/>	men's leotard <input type="checkbox"/>	armsleeve <input type="checkbox"/> L <input type="checkbox"/> R	glove <input type="checkbox"/> L <input type="checkbox"/> R	gauntlet <input type="checkbox"/> L <input type="checkbox"/> R

Donning/Doffing Accessories:

## Adjustable Compression Velcro Wrap: **circaid**<sup>®</sup>



STYLE	<input type="checkbox"/> lower	<input type="checkbox"/> upper	<input type="checkbox"/> foot	<input type="checkbox"/> full leg	<input type="checkbox"/> arm	<input type="checkbox"/> hand	<input type="checkbox"/> nighttime
EXTREMITY	<input type="checkbox"/> left	<input type="checkbox"/> right	<input type="checkbox"/> bilateral				
COMPRESSION	<input type="checkbox"/> 20-30	<input type="checkbox"/> 30-40	<input type="checkbox"/> 40-50				

Check all that apply

\*Dispense as written - No substitution

QTY \_\_\_\_\_ REFILLS \_\_\_\_\_ DURATION \_\_\_\_\_

Patient Name \_\_\_\_\_ DOB \_\_\_\_\_

## Diagnosis:

- 187.2 Venous Insufficiency     197.2 Postmastectomy Lymphedema  
 189.0 Lymphedema     Other Code \_\_\_\_\_

Physician Signature \_\_\_\_\_ Date \_\_\_\_\_

Physician Name \_\_\_\_\_ NPI \_\_\_\_\_

HOME OF THE ORIGINAL  
CASINO PANTY HOSE



Located in the Sahara West Center

**SERVING LAS VEGAS FOR OVER 30 YEARS**

3441 W Sahara Ave. Suite D5  
Las Vegas, NV 89102

[www.mobilemedicalservices.com](http://www.mobilemedicalservices.com)

**(702) 733-1900**

**FAX: 702-737-9354**

Your prescription entitles you to a 10% savings on your vascular stockings.

**MOST INSURANCES ACCEPTED**

## Indications:

### 20–30 mmHg

- For heaviness and fatigue in the legs
- Mild varicosities without significant disposition for edema
- Post sclerotherapy
- Initial varices during pregnancy
- Small varicose dilations of cutaneous veins

### For lymphedema compression of arm

- After mastectomy
- Blood poisoning
- Post traumatic and post-operative edemas
- Tennis elbow syndrome
- Severe burn therapy

### 30–40 mmHg

- Relief of aching heaviness and fatigue caused by varices
- Post sclerotherapy
- Post surgical stripping
- Prophylaxis of thrombosis
- Prophylaxis and treatment of complications of varicose veins and post phlebotic syndrome with chronic venous insufficiency
- Varices during pregnancy
- Pregnant patients with previous phlebitis
- Control of edema and effective scar formation after burns
- Stasis dermatitis due to Chronic Venous Insufficiency (CVI)

### 40–50 mmHg

- For severe degrees of above
- Emphasized edema from above causes
- Correctable lymphedema
- Severe chronic venous insufficiency as with post phlebotic syndrome
- Chronic venous problems after surgery
- Venous ulcers

### Contraindications

- High grade arterial insufficiency
- Wet dermatoses
- Cutaneous infections
- Hyperdermatitis in acute stage